



## Cartilage Restoration Rehabilitation Protocol (Femoral Condyle)

Phase	Goals	Precautions/Restrictions	Treatment
Weeks 0-6	<ul> <li>Protect surgical site</li> <li>Manage swelling and pain</li> <li>Achieve and maintain good quadriceps activation</li> <li>Achieve full ROM by 6 weeks</li> </ul>	<ul> <li>TTWB</li> <li>No ROM restrictions</li> </ul>	<ul> <li>PRICE</li> <li>Quadriceps activation and strength should be emphasized</li> <li>Knee flexion and terminal extension ROM</li> <li>Gentle stretching of hamstrings, calf, quadriceps to tolerance</li> <li>OKC hip strengthening in all planes</li> <li>Ok to initiate stationary biking without resistance</li> <li>Initial visit: FOTO, LEFS</li> </ul>
Weeks 6-12	<ul> <li>Full WB ambulation by week 8-9</li> <li>Ambulate community distances by 12 weeks</li> </ul>	<ul> <li>WBAT progressing to full WB over 2-3 weeks</li> <li>No impact (running, cutting, pivoting)</li> </ul>	<ul> <li>Begin CKC strengthening (avoid anterior knee pain)</li> <li>Normalize calf, hamstring, quadriceps mobility</li> <li>Week 6: FOTO, LEFS</li> </ul>
Weeks 12-24	<ul> <li>Restoring strength of quadriceps, hamstrings, hips</li> <li>Ready to begin impact by 6 months</li> </ul>	<ul> <li>No impact (running, cutting, pivoting)</li> </ul>	<ul> <li>Progress CKC into greater ROM, single leg, multi-planar, and with resistance as tolerated</li> <li>Ok to initiate elliptical</li> <li>Week 12: FOTO, LEFS</li> </ul>
Weeks 24+	<ul> <li>Begin impact training once cleared by MD (jumping, running etc.)</li> </ul>	<ul> <li>Avoid running/jumping on a painful or swollen knee</li> </ul>	<ul> <li>Jumping progression (double to single leg)</li> <li>Return to run program (walk/jog)</li> <li>Anticipated final visit: FOTO, LEFS</li> </ul>

This protocol is not meant to be prescriptive but a recommendation to guide the rehabilitation process. Each patient's progress may vary based on specifics to their injury and procedure.



600 Hennepin Avenue, Minneapolis MN 55403

0

