

SHOULDER ARTHROSCOPY SUBACROMIAL DECOMPRESSION/DISTAL CLAVICLE EXCISION

Rehab Protocol

Weeks 0-4:

- Sling for Comfort (1-2 days) Then discontinue
- Passive to Active shoulder ROM as tolerated
 - 140° Forward Flexion
 - 40° External Rotation with arm at side
 - Internal rotation behind back with gentle posterior capsule stretching
 - No rotation with arm in abduction until 4 wks
- With distal clavicle excision, hold cross body adduction until 8wks.
- Grip Strength, Elbow/Wrist/Hand ROM, Codmans
- Avoid Abduction and 90/90 ER until 8wks

Weeks 4-8:

- Advance ROM as tolerated (Goals FF to 160°, ER to 60°)
 - Begin Isometric exercises
 - Progress deltoid isometrics
 - ER/IR (submaximal) at neutral
- Advance to theraband as tolerated

Weeks 8-12:

- Advance to full, painless ROM
- Continue strengthening as tolerated
- Begin eccentrically resisted motion and closed chain activities
- Only do strengthening 3times/wk to avoid rotator cuff tendonitis



OkorohaPA@gmail.com



600 Hennepin Avenue,
Minneapolis MN 55403



Admin Assistant
612-313-0531